

# Phytochemical analysis of Lablab purpureus L. Seed extracts.

<sup>1</sup>D Sai kumari, <sup>2</sup>Dr Neeti saxena  
<sup>1</sup>Research scholar, <sup>2</sup>Asst.professor  
<sup>1</sup>University college for women

**Abstract** - Phytochemicals are the plant derivatives which are used in mainly in herbal drugs. They cure many endemic, epidemic diseases. In the world only 11% of plants were identified as for mankind. There is a huge need of medicinal plants to be discovered for the day today challenge of world population. In the present study phyto-chemical analysis of Lablab purpureus L. showed and contains sugar, steroids, alkaloids, tannins, flavanoids, saponins, coumarins, terpenoids and other pigments. In the analysis used high water and Methanol solvents were used. This study showed that this phytochemical contents have significant pharmacological activity.

**keywords** - Phytochemical, Lablab purpureus, flavanoid, phenol, solvents.

## Introduction:

*Lablab purpureus L.* Belongs to the family *fabaceae*, commonly called as hyacinth bean, Dolichos bean, Seim bean, Egyptian bean, Kidney bean and Indian bean. Hyacinth is a tropical perennial twinning plant. Flowers appear in clusters with purple or white colour, which develop into flat, thick, slightly curved green pods with pointing ends. In each pod generally six seeds are present, which are mottled, mostly oval in shape. In 100 grams of *Dolichos* bean contain proteins, carbohydrates, dietary fibres, vitamins A, B complex vitamins and minerals like Ca, Fe, Cu, Se, Mg, K and Zn along with bioactive Phytochemical compounds. The present study deals with the phytochemical analysis of *Dolichos* bean !1)

The plant can show the pharmacological effects to the diseases of anti-microbial, antioxidant, anticancer, hypo-lipidemic, central nervous system, cardiovascular, respiratory, immunological, anti-inflammatory, analgesic, anti-pyretic, cytotoxic, insecticidal, anti-diabetic, anti-lithiatic, anemic diseases.

The present study which deals with the phytochemical analysis of *Lablab purpureus L seed* extracts.

**Sample preparation:** Field seed samples collected from the Karimnagar District, Telangana, India. Seeds were washed many times and shadow dried. Seeds were subjected to pulverization to get coarse powder. It was stored in airtight container for further use. Solvents like Methanol and Water were subjected to soxhlet extraction. For every solvent 250mg of seed powder was used against 150 ml of solvent. Fractions were collected after some rounds of successful extraction in the soxhlet apparatus (2).

**Materials and Methods:** All the extracts were used for phytochemical analysis. According to the test of content, test sample varies by following standard protocols and the results were listed in the tabular form

**Test for Reducing sugars:** The extracts were treated with 5.0 ml of Fehling's solution and kept in boiling water bath. The formation of yellow or red colour precipitate indicates the presence of reducing sugars. (3).

**Test for phenols:** 2% FeCl<sub>3</sub> was added to the 1ml of sample formed blue black colouration. This showed positive result for phenols. (4)

**Test for steroids:** 1 ml of the extract was dissolved in 10ml of chloroform and equal volume of concentrated sulphuric acid was added by the sides of the test tube. The upper layer turned red and the sulphuric acid layer formed fluorescent green with yellow. This indicated the presence of steroids. (5).

**Test for Alkaloids:** For the detection of presence of Alkaloid, Wagner's was performed, where initially solvent free extract 50 mg was mixed with few ml of dilute hydrochloric acid and then filtered, the filtrate is used for testing the presence of alkaloids. To a few ml of filtrate, a few drops of Wagner's reagent were added by the side of the test tube. A reddish brown precipitate indicates the presence of alkaloids (6)

**Test for Oil:** A small quantity of extract was pressed between two filter papers; oil stain on the paper indicates the presence of fixed oil [7].

**Test for Tannins:** 10 ml aqueous extract was mixed with 0.1% of Ferric chloride. The positive results for the presence of tannin to form green colour to blue black. (8).

**Test for Flavonoids:** 2 ml of the each extract was added to 2 ml water and 5 ml of 20% NaOH. The yellow colouration formed to show the positive results of flavonoids. (9)

**Test for saponins :** About 2 g of the powdered sample was boiled in 20 ml of distilled water bath and filtered. The 10 ml of the filtrate was mixed with 5 ml of distilled water and shaken vigorously for a suitable persistent froth. The frothing was mixed with 3 drops of olive oil and shaken vigorously, and then the formation of emulsion was observed.(3)

**Test for Coumarins:** 3 ml of 10% NaOH was added to 2ml of aqueous extract. Formation of yellow colour indicated the presence of Coumarins. (5)

**Salkowski's Test:** It is the test for terpenoids. 5ml of each extract was mixed in 2 ml of chloroform and 3ml of concentrated H<sub>2</sub>SO<sub>4</sub> was carefully added to form a layer. A reddish brown colouration of the interface was formed to show positive results for the presence of terpenoid. (8)

**Elemental analysis:** Samples are prepared as by taking 2 gms of samples, and add 10 ml of conc. nitric acid. Transfer them in to crucibles and keep them inside furnace at 200 degrees. Take out the ash formed in the crucibles. Dilute them with water and filter the supernatant. Standards are prepared as per, and readings are noted with the help of Atomic absorption spectroscopy [6].

**Test for carbohydrates:** The extracts were treated with 5.0 ml of Fehling's solution and kept in boiling water bath. The formation of yellow or red colour precipitate indicates the presence of reducing sugars. (3)

**Estimation of crude fiber:** Extract 2 g of ground material of seed with ether or petroleum ether to remove fat (Initial boiling temperature 35-38° C and final temperature 52° C). If fat content is below 1%, extraction may be omitted. After extraction with ether, boil 2 g of dried material with 200 ml of sulphuric acid for 30 min with bumping chips. Filter through muslin and wash with boiling water, until washings are no longer acidic. Boil with 200 ml of sodium hydroxide solution for 30 min. Filter through muslin cloth again and wash with 25 ml of boiling 1.25% H<sub>2</sub>SO<sub>4</sub>, three 50 ml portions of water and 25 ml alcohol. Remove the residue and transfer to ashing dish. Dry the residue for 2 h at 130 ± 2° C. Cool the dish in desiccators. Ignite for 30 min at 600 ± 15° C. Cool in desiccators [6].

**Glycosides:** 50 mg of extract was mixed with few ml of conc. hydrochloric acid for 2 hours on water bath and then filtered; the filtrate was used for testing the presence of glycosides by legal's test. 0.5 ml of filtrate was dissolved in pyridine, and then sodium nitroprusside solution was added and made alkaline using 10% sodium hydroxide. Pink color indicates the presence of glycosides [6].

**Total protein:** The total protein content in the seed S is estimated by Lowry's method. Different dilutions of BSA solutions are prepared by mixing stock BSA solution (1 mg/ml) and water in the test tube, as given in the table. The final volume in each of the test tubes is 5 ml. The BSA range is 0.05-1 mg/ml. From these different dilutions, pipette out 0.2 ml protein solution to different test tubes and add 2 ml of alkaline copper sulphate reagent (analytical reagent). Mix the solutions well. This solution is incubated at room temperature for 10 mins. Then add 0.2 ml of reagent Folin Ciocalteu solution (reagent solutions) to each tube, and incubate for 30 min. Zero the colorimeter with blank and take the optical density (measure the absorbance) at 660 nm. Plot the absorbance against protein concentration to get a standard calibration curve. Check the absorbance of unknown sample, and determine the concentration of the unknown sample using the standard curve plotted above [10]

**Total fat:** The total fat content was determined by extraction of 2.0- 2.5 g of dry ground sample for 12 h in a Soxhlet with petroleum ether, and removed the solvent by rotary evaporator, then dried the sample in hot air oven at 100° C for about 1 h to allow the ether evaporate (11-14).

**Test for Pigments:** 1ml of extract was extracted with 10ml of chloroform in a test tube with vigorous shaking and then 85% sulphuric acid was added. A blue colour at the interface showed the presence of carotenoids. (9)

#### **Test for combined anthraquinones**

1ml of extract was boiled with 2ml of 10% HCl for 5 mins. The cooled filtrate was extracted with equal volume of chloroform and the chloroform layer was transferred to dry clean test tube. 10 % ammonia solution was added in the chloroform layer, shaken well and allowed to separate. The separated layer was observed for colour change. Pink colour is the positive result for combined anthraquinones. (9).

#### **Results and Discussion**

Reducing sugars were tested against Fehling solution showed positive indication of yellow (or) red precipitate appeared in Water and Methanol. Phenol appeared in Water extract where as absent in Methanol, they showed blue and black colouration. Steroids were tested against Chloroform, Positive results appeared green to yellow colouration both in Water & Methanol. Wagner's test was performed to test for Alkaloids appeared as reddish brown precipitate in both in water and methanol extracts. Salkowski's test was done to identify terpenoids in the seeds, reddish brown colour indicates the presence of Alkaloids in water and methanol. They were the important component in the plant which acts as astringent (15). Flavanoids are major group of phytochemicals present in the plants, they were showed yellow colour in both water and methanol. Coumarins are aromatic compounds, widely used to enhance aroma. They are naturally found in plants in a crystalline form, yellow colour formation promised the coumarins presence in the seed, only appeared in water extracts. A clear emulsion was formed in water extract where as absent in methanolic extract. Glycosides present in both extracts showed pink colouration. Carotenoids showed blue colour interface in both extractions. Alkaloids, Fibre, Glycosides and Protein were tested by using standard official methods of AOAC methods. Only fat content was tested by used Petroleum ether.

Phenols, flavonoids tannins and coumarins are responsible for chemo preventive properties like anti-oxidant, anti-carcinogenic and anti-inflammatory effects (16). The Acoholic extracts of *Lablab purpureus* leaves and seeds reduced blood glucose (17), where as methanolic extracts of some doses controls blood glucose ranges as well as body weight in animals(18). Aqueous extract of *Dolichos* bean can increase significant change in heamoglobin percentage (19) which shows anti-inflammatory (20) analgesic effect, antioxidant activity (21) cytotoxic activity (22) Hypolipidemic effect (23) Anti-microbial effect (24) insecticidal effect (25) Hepatoprotective effect (26) Antilithiatic activity (27) along with this side effects and contradictions also present.

### Conclusion

The above study discussed the chemical constituents and pharmacological effects of *Lablab purpureus* have the wide range of medicinal properties have been used for many medical applications .

### Acknowledgement

I express my gratitude to the Department of Botany, University College for Women, University College of Science, Osmania University, Hyderabad, Telangana, India for providing facilities and laboratory

### References:

- [1] "*Lablab purpureus*". Natural Resources Conservation Service PLANTS Database. USDA. Retrieved 22 January 2016.
- [2] Rufus Auxilia, L. et al. Phytochemical analysis of seed extracts *Macrotyloma uniflorum* (horse gram) International Journal of Current Research Vol. 5, Issue, 11, pp.3339-3342, November, 2013 .
- [3] Thenmozhi, A., Nagalakshmi, K., Mahadeva Rao, US. 2011. Qualitative analysis of phytochemicals, and comparative superoxide radical scavenging along with reducing potency of *Solanum nigrum* using various solvent extracts. International journal of Green Pharmacy. 5(4) 318-324.
- [4] Yadhav RNS and Munin Agarwala. 2011 Phytochemical Analysis of some Medicinal Plants. Journal of Phytology. 3(12): 10-14
- [5] Savithramma, N., Linga Rao, M. and Suhurulatha, D. 2011 Screening of Medicinal plant for secondary Metabolites. Middle-East Journal of Scientific Research 8 (3): 579-584.
- [6] AOAC (1995) Official methods of analysis (References: 922.06, 991.20, 923.03 and 978.10). Association of Official Analytical Chemists
- [7] Dangeti SR, Karthikeyan S, Kumar GR, Desai S (2013) Proximate and Phytochemical Analysis of Seed Coat from *P. sumantranse* (Little Millet). Biochem Anal Biochem 2: 134. doi:10.4172/2161-1009.1000134
- [8] .Doss, A. 2009 Preliminary phytochemical screening of some Indian Medicinal Plants. Ancient Science of Life. 29(2) 12-16.
- [9] Moses, A.G. Maobe, Erastus Gatebe, Leonard Gitu and Henry Rotich. 2013 Preliminary Phytochemical Screening of Eight Selected Medicinal Herbs Used for the Treatment of Diabetes, Malaria and Pneumonia in Kisii Region, Southwest Kenya. European Journal of Applied Sciences 5 (1): 01-06.
- [10] Shiva M, Mohammad S, Manoochehr H, Yaghoub A, Seyed ESE, et al. (2007) Antioxidant power of Iranian propolis extract. Food Chem 103: 729-733. 11.
- [11] Paquot C (1979) Standard methods for the analysis of oils, fats and derivatives. (6th Ed.) Part I (Sections I and II) Pure and Appl Chem 51: 2503-2525.
- [12] Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurement with the folin phenol reagent. J Biol Chem 193: 265-275.
- [13] Okezie OB, Bello AB (1988) Physico-chemical and functional properties of winged bean flour and isolate compared with soy isolate. J Food Sci 53: 450- 454.
- [14] Olayanjo TMA (2004) Beniseed: A big treasure in small capsule. Publication in RMRDC. Raw Material Upgrade Magazine, Abuja, Nigeria.
- [15] Rufus Auxilia, L.et.al., Phytochemical anlysis of seed extracts *Macrotyloa uniflorum* (horse gram): International journal of current research vol.5,Issue, 11, pp.3339-3342, November, 2013.
- [16] Huang WY, Cai Yz, Zhang Y. Nutr Cancer.2010:6(2) 1-20. doi : 10.1080/06135580903191585.
- [17] Singh R and Sankar C. Screening for anti-diabetic activity of the ethanolic extract of *Dolichos lablab* leaves. Ph Tech Med 2012; 1(5): 177-180 .
- [18] Ahmed M, Trisha UK, Shaha SR, Dey AK and Rahmatullah M. An initial report on the antihyperglycemic and antinociceptive potential of *Lablab purpureus* L. Sweet whit and purple. IJPSR 2012; 3(3): 776-781.
- [19] Somulung SA, Lucero MA, Niverca MS, Dalin KA, Dejesus R and Domingo ED. In vivostudy on the effect of *Dolichos lablab* (bataw) beans extract against Iron deficiency in *Rattus norvergicus* ( Wistar rat). Faima University Research journal 2012; 4: 112-115.
- [20] Habib MAM, Hasan R, Nayeem J, Uddin N and Rana S. Anti-inflammatory, antioxidant and cytotoxic potential of methanolic extract of two Bangladeshi bean *Lablab purpureus* L. sweet white and purple. IJPSR 2012; 3(3): 776-781
- [21] Teixeira CS, Assreuy AMS, Osterne VJ, Amorim RMF, Brizeno LAC, Debray H, Nagano CS, Delatorre P, Sampaio AH, Rocha BAM and Cavada BS. Mannose-specific legume lectin from the seeds of *Dolichos lablab* (FRIL) stimulates inflammatory and hypernociceptive processes in mice. Process Biochemistry 2014; 49(3): 529–534.
- [22] Nasrin F, Bulbu IJ, Begum Y and Khanum S. In vitro antimicrobial and cytotoxicity screening of nhexane, chloroform and ethyl acetate extracts of *Lablab purpureus* (L.) leaves. Agric Biol J N Am 2012; 3(2): 43-48

- [23] Ramakrishna V, Rani PJ and Rao PR. Hypocholesterolemic effect of diet supplemented with Indian bean (*Dolichos lablab* L. var *lignosus*) seeds. *Nutrition & Food Science* 2007; 37(6): 452 – 456.
- [24] Ye XY, Wang HX and Ng TB. Dolichin, a new chitinase-like antifungal protein isolated from field beans (*Dolichos lablab*). *Biochem Biophys Res Commun* 2000; 269(1): 155-159.
- [25] Janarthanan S, Sakthivelkumar S, Veeramani V, Radhika D and Muthukrishnan S. A new variant of antimetabolic protein, arcelin from an Indian bean, *Lablab purpureus* (Linn.) and its effect on the stored product pest, *Callosobruchus maculatus*. *Food Chem* 2012;135(4):2839-2844
- [26] Im AR, Kim YH, Lee HW and Song KH. Water extract of *Dolichos lablab* attenuates hepatic lipid accumulation in a cellular nonalcoholic fatty liver disease model. *J Med Food* 2016;19(5):495-503.
- [27] Deoda RS, Pandya H, Patel M, Yadav KN, Kadam PV and Patil MJ. Antilithiatic activity of leaves, bulb and stem of *Nymphaea odorata* and *Dolichos lablab* Beans. *Research Journal of Pharmaceutical, Biological and Chemical Sciences* 2012; 3(1): 814- 819.

